

FILE 'HOME' ENTERED AT 17:44:06 ON 15 JUN 2005

=> FIL STNGUIDE

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'STNGUIDE' ENTERED AT 17:44:12 ON 15 JUN 2005

USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE

AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.

LAST RELOADED: Jun 10, 2005 (20050610/UP).

=> FIL HOME

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.06

0.27

FILE 'HOME' ENTERED AT 17:44:16 ON 15 JUN 2005

=> index bioscience

FILE 'DRUGMONOG' ACCESS NOT AUTHORIZED

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.48

INDEX 'ADISCTI, ADISINSIGHT, ADISNEWS, AGRICOLA, ANABSTR, ANTE, AQUALINE, AQUASCI, BIOBUSINESS, BIOCOMMERCE, BIOENG, BIOSIS, BIOTECHABS, BIOTECHDS, BIOTECHNO, CABA, CANCERLIT, CAPLUS, CEABA-VTB, CEN, CIN, CONFSCI, CROPB, CROPU, DDFB, DDFU, DGENE, DISSABS, ...' ENTERED AT 17:44:30 ON 15 JUN 2005

75 FILES IN THE FILE LIST IN STNINDEX

Enter SET DETAIL ON to see search term postings or to view search error messages that display as 0* with SET DETAIL OFF.

=> s byrne m?/au

52 FILE ADISCTI
0* FILE ADISINSIGHT
0* FILE ADISNEWS
59 FILE AGRICOLA
2 FILE ANABSTR
109 FILE ANTE
6 FILE AQUALINE
64 FILE AQUASCI
92 FILE BIOBUSINESS
0* FILE BIOCOMMERCE
25 FILE BIOENG
620 FILE BIOSIS
20 FILE BIOTECHABS
20 FILE BIOTECHDS
112 FILE BIOTECHNO
98 FILE CABA
102 FILE CANCERLIT
328 FILE CAPLUS
13 FILE CEABA-VTB
0* FILE CIN
49 FILE CONFSCI
5 FILE CROPU
16 FILE DDFB

61	FILE DDFU
129	FILE DGENE
53	FILE DISSABS
16	FILE DRUGB
0*	FILE DRUGMONOG2
61	FILE DRUGU
5	FILE EMBAL
432	FILE EMBASE
252	FILE ESBIODBASE
4	FILE FEDRIP
82	FILE FOMAD
0*	FILE FOREGE
170	FILE FROSTI
119	FILE FSTA
7	FILE HEALSAFE
71	FILE IFIPAT
0*	FILE IMSDRUGNEWS
0*	FILE IMSPRODUCT
0*	FILE IMSRESEARCH
5	FILE JICST-EPLUS
152	FILE LIFESCI
0*	FILE MEDICONF
528	FILE MEDLINE
6	FILE NIOSHTIC
25	FILE NTIS
0*	FILE NUTRACEUT
34	FILE OCEAN
379	FILE PASCAL
0*	FILE PCTGEN
56 FILES SEARCHED...	
0*	FILE PHAR
0*	FILE PHARMAML
0*	FILE PHIC
0*	FILE PHIN
224	FILE PROMT
0*	FILE PROUSDDR
0*	FILE RDISCLOSURE
913	FILE SCISEARCH
208	FILE TOXCENTER
64	FILE USPATFULL
4	FILE USPAT2
1	FILE VETB
3	FILE WATER
88	FILE WPIDS
88	FILE WPINDEX

49 FILES HAVE ONE OR MORE ANSWERS, 75 FILES SEARCHED IN STNINDEX

L1 QUE BYRNE M?/AU

=> s goke b?/au

23	FILE ADISCTI
0*	FILE ADISINSIGHT
0*	FILE ADISNEWS
5	FILE AGRICOLA
0*	FILE BIOCOMMERCE
99	FILE BIOSIS
1	FILE BIOTECHABS
1	FILE BIOTECHDS
66	FILE BIOTECHNO
13	FILE CABA
57	FILE CANCERLIT
61	FILE CAPLUS
0*	FILE CIN

21 FILE DDFU
 12 FILE DGENE
 0* FILE DRUGMONOG2
 21 FILE DRUGU
 2 FILE EMBAL
 280 FILE EMBASE
 120 FILE ESBIODBASE
 0* FILE FOREGE

39 FILES SEARCHED...

1 FILE IFIPAT
 0* FILE IMSDRUGNEWS
 0* FILE IMSPRODUCT
 0* FILE IMSRESEARCH
 1 FILE LIFESCI
 0* FILE MEDICONF
 268 FILE MEDLINE
 0* FILE NUTRACEUT
 16 FILE PASCAL
 0* FILE PCTGEN
 0* FILE PHAR
 0* FILE PHARMAML
 0* FILE PHIC
 0* FILE PHIN
 0* FILE PROUSDDR
 0* FILE RDISCLOSURE
 452 FILE SCISEARCH
 62 FILE TOXCENTER
 1 FILE USPATFULL
 4 FILE WPIDS
 4 FILE WPINDEX

24 FILES HAVE ONE OR MORE ANSWERS, 75 FILES SEARCHED IN STNINDEX

L2 QUE GOKE B?/AU

=> s 11 and 12

2 FILE ADISCTI
 0* FILE ADISINSIGHT
 0* FILE ADISNEWS
 1 FILE AGRICOLA
 0* FILE BIOCOMMERCE
 4 FILE BIOSIS
 1 FILE BIOTECHNO
 1 FILE CANCERLIT
 4 FILE CAPLUS
 0* FILE CIN
 2 FILE DDFU
 6 FILE DGENE
 0* FILE DRUGMONOG2
 2 FILE DRUGU
 7 FILE EMBASE
 6 FILE ESBIODBASE
 0* FILE FOREGE
 0* FILE IMSDRUGNEWS
 0* FILE IMSPRODUCT
 0* FILE IMSRESEARCH
 0* FILE MEDICONF

49 FILES SEARCHED...

6 FILE MEDLINE
 0* FILE NUTRACEUT
 0* FILE PCTGEN
 0* FILE PHAR
 0* FILE PHARMAML
 0* FILE PHIC

0* FILE PHIN
 0* FILE PROUSDDR
 0* FILE RDISCLOSURE
 13 FILE SCISEARCH
 1 FILE TOXCENTER
 1 FILE WPIDS
 1 FILE WPINDEX

16 FILES HAVE ONE OR MORE ANSWERS, 75 FILES SEARCHED IN STNINDEX

L3 QUE L1 AND L2

=> file hits

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
2.95	3.43

FULL ESTIMATED COST

FILE 'SCISEARCH' ENTERED AT 17:47:20 ON 15 JUN 2005

Copyright (c) 2005 The Thomson Corporation

FILE 'EMBASE' ENTERED AT 17:47:20 ON 15 JUN 2005

COPYRIGHT (C) 2005 Elsevier Inc. All rights reserved.

FILE 'DGENE' ENTERED AT 17:47:20 ON 15 JUN 2005

COPYRIGHT (C) 2005 THE THOMSON CORPORATION

FILE 'ESBIOBASE' ENTERED AT 17:47:20 ON 15 JUN 2005

COPYRIGHT (C) 2005 Elsevier Science B.V., Amsterdam. All rights reserved.

FILE 'MEDLINE' ENTERED AT 17:47:20 ON 15 JUN 2005

FILE 'BIOSIS' ENTERED AT 17:47:20 ON 15 JUN 2005

Copyright (c) 2005 The Thomson Corporation

FILE 'CAPLUS' ENTERED AT 17:47:20 ON 15 JUN 2005

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'ADISCTI' ENTERED AT 17:47:20 ON 15 JUN 2005

COPYRIGHT (C) 2005 Adis Data Information BV

FILE 'DDFU' ACCESS NOT AUTHORIZED

FILE 'DRUGU' ENTERED AT 17:47:20 ON 15 JUN 2005

COPYRIGHT (C) 2005 THE THOMSON CORPORATION

FILE 'AGRICOLA' ENTERED AT 17:47:20 ON 15 JUN 2005

FILE 'BIOTECHNO' ENTERED AT 17:47:20 ON 15 JUN 2005

COPYRIGHT (C) 2005 Elsevier Science B.V., Amsterdam. All rights reserved.

FILE 'CANCERLIT' ENTERED AT 17:47:20 ON 15 JUN 2005

FILE 'TOXCENTER' ENTERED AT 17:47:20 ON 15 JUN 2005

COPYRIGHT (C) 2005 ACS

FILE 'WPIDS' ENTERED AT 17:47:20 ON 15 JUN 2005

COPYRIGHT (C) 2005 THE THOMSON CORPORATION

FILE 'WPINDEX' ACCESS NOT AUTHORIZED

=> s 13

L4 55 L3

=> dup rem l4
DUPLICATE IS NOT AVAILABLE IN 'DGENE'.
ANSWERS FROM THESE FILES WILL BE CONSIDERED UNIQUE
PROCESSING COMPLETED FOR L4

L5 29 DUP REM L4 (26 DUPLICATES REMOVED)
ANSWERS '1-13' FROM FILE SCISEARCH
ANSWERS '14-16' FROM FILE EMBASE
ANSWERS '17-22' FROM FILE DGENE
ANSWERS '23-24' FROM FILE ESBIODBASE
ANSWERS '25-26' FROM FILE BIOSIS
ANSWER '27' FROM FILE CAPLUS
ANSWER '28' FROM FILE ADISCTI
ANSWER '29' FROM FILE DRUGU

=> s l5 and py<1999
4 FILES SEARCHED...
6 FILES SEARCHED...
10 FILES SEARCHED...
13 FILES SEARCHED...
L6 9 L5 AND PY<1999

=> d bib abs 1-9

L6 ANSWER 1 OF 9 SCISEARCH COPYRIGHT (c) 2005 The Thomson Corporation on
STN
AN 1998:606006 SCISEARCH
GA The Genuine Article (R) Number: 106XW
TI Glucagon-like peptide 1 improves the ability of the beta-cell to sense and
respond to glucose in subjects with impaired glucose tolerance
AU **Byrne M M (Reprint);** Gliem K; Wank U; Arnold R; Katschinski M;
Polonsky K S; **Goke B**
CS UNIV MARBURG, DEPT INTERNAL MED, CLIN RES UNIT GASTROINTESTINAL
ENDOCRINOL, D-35033 MARBURG, GERMANY (Reprint); UNIV CHICAGO, DEPT MED,
CHICAGO, IL 60637; PRITZKER SCH MED, CHICAGO, IL
CYA GERMANY; USA
SO DIABETES, (AUG 1998) Vol. 47, No. 8, pp. 1259-1265.
Publisher: AMER DIABETES ASSOC, 1660 DUKE ST, ALEXANDRIA, VA 22314.
ISSN: 0012-1797.
DT Article; Journal
FS LIFE; CLIN
LA English
REC Reference Count: 46
ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS
AB Impaired glucose tolerance (IGT) and NIDDM are both associated with an
impaired ability of the P-cell to sense and respond to small changes in
plasma glucose concentrations. The aim of this study was to establish if
glucagon-like peptide 1 (GLP-1), a natural enteric peptide and potent
insulin secretagogue, improves this defect. Two weight-matched groups, one
with eight subjects having IGT (2-h glucose, 10.1 +/- 0.3 mmol/l) and
another with seven subjects with diet-treated NIDDM (2-h glucose, 14.5 +/-
0.9 mmol/l), were studied on two occasions during a 12-h oscillatory
glucose infusion, a sensitive test of the ability of the beta-cell to
sense and respond to glucose. Glucose was infused with a mean rate of 4 mg
. kg(-1). min(-1), amplitude 33% above and below the mean rate, and
periodicity of 144 min, with infusion of saline or GLP-1 at 0.4 pmol .
kg(-1). min(-1) for 12 h. Mean glucose levels were significantly lower in
both groups during the GLP-1 infusion compared with during saline
infusion: 9.2 +/- 0.4 vs. 6.4 +/- 0.1 mmol/l in the IGT subjects (P <
0.0004) and 14.6 +/- 1.0 vs. 9.3 +/- 0.7 mmol/l in NIDDM subjects (P <
0.0002). Despite this significant reduction in plasma glucose
concentration, insulin secretion rates (ISRs) increased significantly in
IGT subjects (513.3 +/- 77.6 vs. 583.1 +/- 100.7 pmol/min; P < 0.03), with
a trend toward increasing in NIDDM subjects (561.7 +/- 122.16 vs. 642.8

+/- 128 pmol/min; P = 0.1). These results were compatible with enhanced insulin secretion in the presence of GLP-1. Spectral power was used as a measure of the ability of the P-cell to secrete insulin in response to small changes in the plasma glucose concentration during the oscillatory infusion. Spectral power for ISR increased from 2.1 +/- 0.9 during saline infusion to 7.4 +/- 1.3 during GLP-1 infusion in IGT subjects (P < 0.004), but was unchanged in NIDDM subjects (1.0 +/- 0.4 to 1.5 +/- 0.6; P = 0.3). We concluded that low dosage GLP-1 improves the ability of the beta-cell to secrete insulin in both IGT and NIDDM subjects, but that the ability to sense and respond to subtle changes in plasma glucose is improved in IGT subjects, with only a variable response in NIDDM subjects. beta-Cell dysfunction was improved by GLP-1 infusion, suggesting that early GLP-1 therapy may preserve beta-cell function in subjects with IGT or mild NIDDM.

L6 ANSWER 2 OF 9 SCISEARCH COPYRIGHT (c) 2005 The Thomson Corporation on STN
 AN 1998:462137 SCISEARCH
 GA The Genuine Article (R) Number: ZL335
 TI GLP-1 improves first phase insulin secretion without altering insulin sensitivity in subjects with impaired glucose tolerance
 AU **Byrne M (Reprint)**; Ulrich W; Katschinski M; **Goke B**
 SO DIABETES, (MAY 1998) Vol. 47, Supp. [1], pp. 744-744.
 Publisher: AMER DIABETES ASSOC, 1660 DUKE ST, ALEXANDRIA, VA 22314.
 ISSN: 0012-1797.
 DT Conference; Journal
 FS LIFE; CLIN
 LA English
 REC Reference Count: 0

L6 ANSWER 3 OF 9 SCISEARCH COPYRIGHT (c) 2005 The Thomson Corporation on STN
 AN 1998:191995 SCISEARCH
 GA The Genuine Article (R) Number: YZ534
 TI Inhibitory effects of hyperglycaemia on fed jejunal motility: potential role of hyperinsulinaemia
 AU **Byrne M M (Reprint)**; Pluntke K; Wank U; Schirra J; Arnold R; **Goke B**; Katschinski M
 CS UNIV MARBURG, DEPT GASTROENTEROL & ENDOCRINOL, CLIN RES UNIT GASTROINTESTINAL ENDOCRINOL, D-35033 MARBURG, GERMANY (Reprint)
 CYA GERMANY
 SO EUROPEAN JOURNAL OF CLINICAL INVESTIGATION, (JAN 1998) Vol. 28, No. 1, pp. 72-78.
 Publisher: BLACKWELL SCIENCE LTD, P O BOX 88, OSNEY MEAD, OXFORD, OXON, ENGLAND OX2 ONE.
 ISSN: 0014-2972.
 DT Article; Journal
 FS LIFE; CLIN
 LA English
 REC Reference Count: 33

ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS

AB Background Acute hyperglycaemia is known to inhibit jejunal interdigestive motility. This study was undertaken to establish the effects of hyperglycaemia on fed jejunal motility and small intestinal transit time, and to establish if the effects of hyperglycaemia are mediated in part by hyperinsulinaemia.

Methods Nine healthy male volunteers were studied in random order using three experimental conditions: (a) euglycaemic clamp (glucose 5 mmol L⁻¹); (b) hyperglycaemic clamp (glucose 15 mmol L⁻¹); and (c) euglycaemic hyperinsulinaemic clamp (glucose 5 mmol L⁻¹). Fed jejunal motility was induced by an intrajejunal perfusion of lipid (Lipofundin medium-chained triglyceride 10%) at 1.5 mL min⁻¹ (1.5 kcal min⁻¹) for 180 min through the most proximal port of a manometry catheter (eight ports spaced at 2-cm intervals) located just distal to the ligament of Treitz. One minute after

starting the lipid perfusion, 15 g of lactulose dissolved in 20 mL of tap water was infused. Small intestinal transit time was measured by the hydrogen breath test.

Results Acute hyperglycaemia reduced the total number of jejunal contractions and progradely propagated contractions, the motility index ($P < 0.05$) and the mean amplitude of contractions and delayed intestinal transit time. Hyperinsulinaemia reduced the total number of jejunal contractions, motility index ($P < 0.05$) and intestinal transit time.

Conclusions Thus, hyperinsulinaemia may contribute to the inhibitory effects of hyperglycaemia on jejunal motility. In addition, this study demonstrated that intrajejunal infusion of lipid stimulates sustained glucagon-like peptide-1 release. In contrast to fat-induced gastric inhibitory polypeptide release, this glucagon-like peptide-1 release is not inhibited by exogenous or endogenous hyperinsulinaemia ($P = 0.59$).

L6 ANSWER 4 OF 9 SCISEARCH COPYRIGHT (c) 2005 The Thomson Corporation on
STN
AN 97:412985 SCISEARCH
GA The Genuine Article (R) Number: WX380
TI Glucagon-like peptide-1 improves the ability of the beta-cell to sense and
respond to glucose in subjects with impaired glucose tolerance.
AU **Byrne M (Reprint); Kliem K; Wank U; Katschinski M; Arnold R;**
Polonsky K; Goke B
SO DIABETES, (MAY 1997) Vol. 46, Supp. [1], pp. 127-127.
Publisher: AMER DIABETES ASSOC, 1660 DUKE ST, ALEXANDRIA, VA 22314.
ISSN: 0012-1797.
DT Conference; Journal
FS LIFE; CLIN
LA English
REC Reference Count: 0

L6 ANSWER 5 OF 9 SCISEARCH COPYRIGHT (c) 2005 The Thomson Corporation on
STN
AN 96:790333 SCISEARCH
GA The Genuine Article (R) Number: VN947
TI HUMAN STUDIES WITH GLUCAGON-LIKE-PEPTIDE-1 - POTENTIAL OF THE GUT HORMONE /
FOR CLINICAL USE
AU **BYRNE M M (Reprint); GOKE B**
CS UNIV MARBURG, DEPT INTERNAL MED, CLIN RES UNIT GASTROINTESTINAL
ENDOCRINOL, D-3550 MARBURG, GERMANY (Reprint)
CYA GERMANY
SO DIABETIC MEDICINE, (OCT 1996) Vol. 13, No. 10, pp. 854-860.
ISSN: 0742-3071.
DT General Review; Journal
FS CLIN
LA ENGLISH
REC Reference Count: 69

ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS

AB So far, a wealth of data originating from in vitro or animal experiments has been collected supporting the concept that the gut hormone, glucagon-like peptide-1 (GLP-1) may serve as a model molecule for the design of a new drug for the treatment of diabetes mellitus. This is supported by observations that GLP-1 has potent insulinotropic action in patients with non-insulin-dependent diabetes mellitus (NIDDM). It enhances beta-cell sensitivity to glucose stimulated insulin secretion. GLP-1 may also have a role in the treatment of impaired glucose tolerance, where the beta-cell is already insensitive to changes in plasma glucose concentrations. It may, as has previously been shown in animal models of 'prediabetes', delay the progressive decline in glucose tolerance to NIDDM. The glucose-dependent action of this peptide is an important feature in the treatment of NIDDM as it will protect against hypoglycaemic reactions, the most serious acute side-effect of antidiabetic therapy. Glucose utilization may be enhanced which would improve metabolic control in both NIDDM and IDDM. A glucagon lowering effect will further enhance

metabolic control. This article reviews current experiences of the effects of GLP-1 in human studies. It points out the outcomes and limitations of previous trials and discusses future directions for the investigation of its potential use as a new agent in diabetes treatment.

L6 ANSWER 6 OF 9 SCISEARCH COPYRIGHT (c) 2005 The Thomson Corporation on STN
AN 96:593240 SCISEARCH
GA The Genuine Article (R) Number: VA493
TI INHIBITORY EFFECTS OF HYPERGLYCEMIA AND HYPERINSULINEMIA ON POSTPRANDIAL HUMAN JEJUNAL MOTILITY
AU **BYRNE M M (Reprint)**; PLUNTKE K; ARNOLD R; **GOKE B**; SCHIRRA J; KATSCHINSKI M
CS UNIV MARBURG, DEPT GASTROINTESTINAL ENDOCRINOL, D-3550 MARBURG, GERMANY
CYA GERMANY
SO DIABETOLOGIA, (AUG 1996) Vol. 39, Supp. 1, pp. 592.
ISSN: 0012-186X.
DT Conference; Journal
FS LIFE; CLIN
LA ENGLISH
REC No References

L6 ANSWER 7 OF 9 SCISEARCH COPYRIGHT (c) 2005 The Thomson Corporation on STN
AN 96:336245 SCISEARCH
GA The Genuine Article (R) Number: UF737
TI INHIBITORY EFFECTS OF HYPERGLYCEMIA AND HYPERINSULINEMIA ON POSTPRANDIAL HUMAN JEJUNAL MOTILITY
AU **BYRNE M (Reprint)**; PLUNTKE K; WANK U; EHLENZ K; **GOKE B**; SCHIRRA J; KATSCHINSKI M
CS UNIV MARBURG, DEPT GASTROENTEROL, W-3550 MARBURG, GERMANY
CYA GERMANY
SO GASTROENTEROLOGY, (APR 1996) Vol. 110, No. 4, Supp. S, pp. A1061.
ISSN: 0016-5085.
DT Conference; Journal
FS LIFE; CLIN
LA ENGLISH
REC No References

L6 ANSWER 8 OF 9 BIOSIS COPYRIGHT (c) 2005 The Thomson Corporation on STN
AN 1996:451839 BIOSIS
DN PREV199699174195
TI Inhibitory effects of hyperglycemia and hyperinsulinemia on postprandial human jejunal motility.
AU **Byrne, M. M.**; Pluntke, K.; Arnold, R.; **Goke, B.**; Schirra, J.; Katschinski, M.
CS Dep. Gastrointestinal Endocrinology, Univ. Marburg, Marburg, Germany
SO Diabetologia, (1996) Vol. 39, No. SUPPL. 1, pp. A156.
Meeting Info.: 32nd Annual Meeting of the European Association for the Study of Diabetes. Vienna, Austria. September 1-5, 1996.
CODEN: DBTGAI. ISSN: 0012-186X.
DT Conference; (Meeting)
Conference; Abstract; (Meeting Abstract)
Conference; (Meeting Poster)
LA English
ED Entered STN: 7 Oct 1996
Last Updated on STN: 7 Oct 1996

L6 ANSWER 9 OF 9 DRUGU COPYRIGHT 2005 THE THOMSON CORP on STN
AN 1998-45051 DRUGU T E
TI GLP-1 improves first phase insulin secretion without affecting insulin sensitivity in subjects with impaired glucose tolerance.
AU **Byrne M**; Ulrich W; Katschinski M; **Goke B**

LO Marburg, Ger.
 SO Diabetes (47, Suppl. 1, A192, 1998)
 CODEN: DIAEAZ ISSN: 0012-1797
 AV No Reprint Address.T
 LA English
 DT Journal
 FA AB; LA; CT
 FS Literature
 AN 1998-45051 DRUGU T E
 AB I.v. infusion of glucagon-like peptide I (GLP-I) 0.4 pmol/kg/min for 30 min increased the acute insulin response to an i.v. glucose tolerance test, compared with saline infusion, 173.7 vs. 98.1 pmol/l/min, without affecting insulin sensitivity or glucose effectiveness, in 6 subjects (mean age 52 yr) with impaired glucose tolerance or early untreated non-insulin dependent diabetes. It is concluded that low-dose GLP-I infusion improves 1st phase insulin secretion in response to i.v. glucose. (conference abstract). (No EX).
 ABEX (E33/JB)

=> log y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

83.09

86.52

STN INTERNATIONAL LOGOFF AT 17:53:41 ON 15 JUN 2005